

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/063194 A1

(51) International Patent Classification⁷: C07D 417/12,
A61K 31/427, A61P 3/10

(21) International Application Number:
PCT/US2003/037089

(22) International Filing Date:
16 December 2003 (16.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/438,538 6 January 2003 (06.01.2003) US

(71) Applicant (for all designated States except US): ELI
LILLY AND COMPANY [US/US]; Lilly Corporate
Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): WEICHERT,
Andreas, Gerhard [DE/DE]; Lilly Forschung GmbH,
Essener Strasse 93, 22419 Hamburg (DE). BARRETT,
David, Gene [US/DE]; Lilly Forschung GmbH, Essener
Strasse 93, 22419 Hamburg (DE). HEUSER, Stefan
[DE/DE]; Lilly Forschung GmbH, Essener Strasse 93,
22419 Hamburg (DE). RIEDL, Rainer [DE/DE]; Lilly

Forschung GmbH, Essener Strasse 93, 22419 Hamburg
(DE). TEBBE, Mark, Joseph [US/US]; 608 Oxboro
Circle, Durham, NC 27713 (US). ZALIANI, Andrea
[IT/DE]; Lilly Forschung GmbH, Essener Strasse 93,
22419 Hamburg (DE).





(74) Agents: WELCH, Lawrence, T. et al.; ELI LILLY AND
COMPANY, P. O. Box 6288, Indianapolis, IN 46206-6288
(US).

(81) Designated States (national): AE, AG, AI., AM, AT (util-
ity model), AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE
(utility model), DE, DK (utility model), DK, DM, DZ, EC,
EE (utility model), EE, EG, ES, FI (utility model), FI, GB,
GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL,
PT, RO, RU, SC, SD, SE, SG, SK (utility model), SK, SI.,
SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: HETEROARYL COMPOUNDS



or



(57) Abstract: According to the present invention there is provided a compound of formula (I) wherein Z is R³ is lower alkyl or halo lower alkyl having from 2 to 6 carbon atoms or arylalkyl or -(CH₂)_s-V where V is a 3 to 8-membered ring which is cycloalkyl, cycloalkenyl, or heterocycloalkyl having one heteroatom selected from oxygen and sulfur; s is independently 0, 1 or 2; M is hydrogen or halo or lower alkyl or perfluoro-lower alkyl; A is or where Y is oxygen or sulfur, and its pharmaceutically acceptable salts thereof. These compounds are considered to be useful for the treatment of type II Diabetes.

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR,

BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

-1-

HETEROARYL COMPOUNDS

Glucokinase (GK, Hexokinase IV) is one of four hexokinases that are found in mammals [Colowick, S.P., in *The Enzymes*, Vol. 9 (P. Boyer, ed.) Academic Press, New York, NY, pages 1-48, 1973]. The hexokinases catalyze the first step in the metabolism of glucose, i.e., the conversion of glucose to glucose-6-phosphate, Glucokinase has a limited cellular distribution, being found principally in pancreatic beta-cells and hepatocytes. In addition, GK is a rate-controlling enzyme for glucose metabolism in these two cell types that are known to play critical roles in whole-body glucose homeostasis [Chipkin, S.R., Kelly, K.L., and Ruderman, N.B. in *Joslin's Diabetes* (C.R. Khan and G.C. Wier, eds.), Lea and Febiger, Philadelphia, PA, pages 97-115, 1994]. The concentration of glucose at which GK demonstrates half-maximal activity is approximately 8 mM. The other three hexokinases are saturated with glucose at much lower concentrations (<1 mM). Therefore, the flux of glucose through the GK pathway rises as the concentration of glucose in the blood increases from fasting (5 mM) to postprandial (\approx 10-15 mM) levels following a carbohydrate-containing meal [Printz, R.G., Magnuson, M.A., and Granner, D.K. in *Ann. Rev. Nutrition* Vol. 13 (R.E. Olson, D.M. Bier, and D.B. McCormick, eds.), Annual Review, Inc., Palo Alto, CA, pages 463-496, 1993]. These findings contributed over a decade ago to the hypothesis that GK functions as a glucose sensor in beta-cells and hepatocytes (Meglasson, M.D. and Matschinsky, F.M. *Amer. J. Physiol.* 246, E1-E13, 1984). In recent years, studies in transgenic animals have confirmed that GK plays a critical role in whole-body glucose homeostasis. Animals that do not express GK die within days of birth with severe diabetes while animals overexpressing GK have improved glucose tolerance (Grupe, A., Hultgren, B., Ryan, A. et al., *Cell* 83, 69-78, 1995; Ferrie, T., Riu, E., Bosch, F. et al., *FASEB J.*, 10, 1213-1218, 1996). An increase in glucose exposure is coupled through GK in beta-cells to increased insulin secretion and in hepatocytes to increased glycogen deposition.


The finding that type II maturity-onset diabetes of the young (MODY-2) is caused by loss of function mutations in the GK gene suggests that GK also functions as a glucose sensor in humans (Liang, Y., Kesavan, P., Wang, L. et al., *Biochem. J.* 309, 167-173, 1995). Additional evidence supporting an important role for GK in the regulation of glucose metabolism in humans was provided by the identification of patients that express

-2-

a mutant form of GK with increased enzymatic activity. These patients exhibit a fasting hypoglycemia associated with an inappropriately elevated level of plasma insulin (Glaser, B., Kesavan, P., Heyman, M. et al., *New England J. Med.* 338, 226-230, 1998). While mutations of the GK gene are not found in the majority of patients with type II diabetes,


- 5 compounds that activate GK and, thereby, increase the sensitivity of the GK sensor system will still be useful in the treatment of the hyperglycemia characteristic of all type II diabetes. Glucokinase activators will increase the flux of glucose metabolism in beta-cells and hepatocytes, which will be coupled to increased insulin secretion and increased glucose utilization and glycogen synthesis. Such agents would be useful for treating type
- 10 II diabetes.

According to the present invention there is provided compounds of formula (I):



wherein Z is

15




R^3 is lower alkyl or halo lower alkyl having from 2 to 6 carbon atoms or arylalkyl or $-(\text{CH}_2)_s\text{-V}$ where V is a 3 to 8-membered ring which is cycloalkyl, cycloalkenyl, or

- 20 heterocycloalkyl having one heteroatom selected from oxygen and sulfur;

s is independently 0, 1 or 2;

M is hydrogen or halo or lower alkyl or perfluoro-lower alkyl;

A is



25

-3-

where Y is oxygen or sulfur, and

R¹ and R² are independently hydrogen, halo, amino, hydroxyamino, nitro, cyano, sulfonamido, lower alkyl, -OR⁵, -COOR⁵, perfluoro- lower alkyl, lower alkyl thio, perfluoro-lower alkyl thio, lower alkyl sulfonyl, perfluoro-lower alkyl sulfonyl, lower alkyl sulfinyl,

R⁵ is hydrogen, lower alkyl or perfluoro-lower alkyl; or furthermore

R² can be -(CH₂)_n-NR⁶R⁷, with n=1,2,3 or 4, and

R⁶ and R⁷ are independently hydrogen or lower alkyl; or together with the nitrogen atom to which they are attached form a five or six-membered heteroaromatic ring containing from 1 to 3 heteroatoms selected from sulfur, oxygen or nitrogen; or a saturated 5- or 6-membered cycloheteroalkyl ring, which contains from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen; or

R² can be alkynyl, substituted with hydrogen, lower alkyl, hydroxy lower alkyl, lower alkoxy lower alkyl, an unsubstituted or hydroxy substituted cycloalkyl ring containing 5 or 6 carbon atoms, a five- or six-membered saturated heterocyclic ring which contains from 1 to 3 hetero atoms selected from the group consisting of sulfur, oxygen or nitrogen, or an unsubstituted five- or six-membered heteroaromatic ring, connected by a ring carbon atom, which contains from to 3 heteroatoms in the ring selected from the group consisting of sulfur, nitrogen and oxygen, or -(CH₂)_n-NR⁸R⁹, with n=1,2, and

R⁸ and R⁹ are independently hydrogen or lower alkyl; or together with the nitrogen atom to which they are attached form a five or six-membered heteroaromatic ring containing from 1 to 3 heteroatoms selected from sulfur, oxygen or nitrogen; or a saturated 5- or 6-membered cycloheteroalkyl ring, which contains from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen; or

R² can be R¹⁰-[(CH₂)_y-W]_z-, with

W is oxygen, sulfur, -SO-, -SO₂-, and

R¹⁰ is a heteroaromatic ring, connected by a ring carbon atom, which contains from 5 to 6 ring members with from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur or nitrogen, or

aryl containing 6 or 10 ring carbon atoms, or

-4-

- aryl containing 6 ring carbon atoms fused with a heteroaromatic ring containing 5 or 6 ring members with 1 or 2 heteroatoms in the ring being selected from the group consisting of nitrogen, oxygen or sulfur, or
- a saturated 5- or 6-membered cycloheteroalkyl ring, which contains from 1 to 2
- 5 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen, or
- a cycloalkyl ring having 5 or 6 carbon atoms, or
- NR¹¹R¹², with R¹¹ and R¹² being independently hydrogen or lower alkyl;
- y is independently 0,1,2,3 or 4; z is independently 0,1; or
- R² can be R¹³-(CH₂)_t-U-, with
- 10 U is -NHCO-, -CONH-, -NHSO₂-, -SO₂NH- and
- R¹³ in the same meaning of R¹⁰ and
- perfluoro-lower alkyl, lower alkyl, lower alkoxy carbonyl or
- NR¹⁴R¹⁵, R¹⁴ and R¹⁵ are independently hydrogen or lower alkyl; or together with the
- nitrogen atom to which they are attached form a five or six-membered heteroaromatic ring
- 15 containing from 1 to 3 heteroatoms selected from sulfur, oxygen or nitrogen; or a saturated
- 5- or 6- membered heterocycloalkyl ring, which contains from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen;
- t is an integer being 0, 1, 2, 3 or 4;
- R⁴ is -C(O)NHR¹⁶, or is R¹⁷;
- 20 R¹⁶ is hydrogen, lower alkyl, lower alkenyl, hydroxy lower alkyl,
- (CH₂)_n-COOR¹⁸, -CO-(CH₂)_n-COOR¹⁹;
- R¹⁷ is an unsubstituted, mono- or di-substituted five- or six-membered heteroaromatic ring
- connected by a ring carbon atom to the amide group shown, which five- or six-membered
- heteroaromatic ring contains from 1 to 4 heteroatoms selected from sulfur, oxygen or
- 25 nitrogen, with one heteroatom being nitrogen which is adjacent to the connecting ring
- carbon atom; said mono- or di-substituted heteroaromatic ring being mono- or di-
- substituted at a position on a ring carbon atom other than adjacent to said connecting
- carbon atom with a substituent selected from the group consisting of lower alkyl, halo,
- nitro, cyano, -(CH₂)_n-OR²⁰, -(CH₂)_n-COOR²¹,
- 30 -(CH₂)_n-CONHR²², -(CH₂)_n-NHR²³,
- n is 0, 1, 2, 3 or 4;
- R¹⁸, R¹⁹, R²⁰, R²¹, R²² and R²³ are independently hydrogen or lower alkyl,

-5-

and its pharmaceutically acceptable salts thereof.

The compounds of formula I have been found to activate glucokinase *in vitro*. Preferably the compounds of formula I have an enhanced solubility profile and further, have improved metabolic stability over the compounds of the prior art. They are
5 particularly useful for increasing insulin secretion in the treatment of type II diabetes.

The present invention provides pharmaceutical compositions comprising a compound of formula I, or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable diluent or carrier.

Further, the present invention provides a compound of formula I, or a
10 pharmaceutically acceptable salt thereof for use as a pharmaceutical; and a compound of formula I, or a pharmaceutically acceptable salt thereof for use in the treatment or prophylaxis of type II diabetes.

The present invention also provides the use of a compound of formula I, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the
15 treatment or prophylaxis of disorders associated with GK dysfunction in mammals; the use of a compound of formula I, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment or prophylaxis of type II diabetes.

The present invention further relates to processes for the preparation of the compounds of formula I, or pharmaceutically acceptable salts thereof. In addition, the
20 present invention relates to a method for the prophylactic or therapeutic treatment of type II diabetes, which method comprises administering an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof; to a human being or an animal in need thereof.


The present invention includes the pharmaceutically acceptable salts of the
25 compounds of formula I. As used herein, the term "pharmaceutically acceptable salts" include acid addition salts, including salts formed with inorganic acids, for example hydrochloric, hydrobromic, nitric, sulphuric or phosphoric acids, or with organic acids, such as organic carboxylic or organic sulphonic acids, for example, acetoxybenzoic, citric, glycolic, *o*- mandelic-l, mandelic-dl, mandelic d, maleic, mesotartaric monohydrate,
30 hydroxymaleic, fumaric, lactobionic, malic, methanesulphonic, napsylic, naphthalenedisulfonic, naphtoic, oxalic, palmitic, phenylacetic, propionic, pyridyl hydroxy pyruvic, salicylic, stearic, succinic, sulfanilic, tartaric, 2-hydroxyethane sulphonic,

-6-

toluene-p-sulphonic, and xinafoic acids. The term "pharmaceutically acceptable salts" also includes any pharmaceutically acceptable base salt such as amine salts, trialkyl amine salts and the like.

In addition to the pharmaceutically acceptable salts, other salts are included in the invention. They may serve as intermediates in the purification of compounds or in the preparation of other, for example pharmaceutically acceptable, acid addition salts, or are useful for identification, characterisation or purification.


It will be appreciated that the compounds of the invention can contain one or more asymmetric carbon atoms which gives rise to isomers. In the compound of formula I the "*" designates an asymmetric carbon atom in this compound. Preferably, the compound of formula I has the following spatial orientation:



wherein, A, M, R^3 and R^4 are as defined above. It will be appreciated that the specific configuration of the carbon to which A is attached, namely "R" or "S", will depend upon the definition of R^3 , M and Y and their respective priorities according to the Cahn-Ingold-Prelog sequence rules as described in J. March, Fourth Edition, Chapter 4, page 109.

Where M is other than hydrogen an additional chiral center at the connecting carbon atom is generated. At this centre the compounds of formula I may be present as a racemate or in the "R" or "S" configuration. Racemic mixtures, enantiomers, diastereomers and individual isomers form part of the present invention, the compounds being employed as racemates or in enantiomerically pure form.

In accordance with this invention, for compounds of formula I where Z is



"Δ" denotes compounds wherein "A" and "M" have a trans configuration across the double bond.

-7-

The term "trans" as utilized in this application, designates that the two substituents attached at each end of the double bond are on opposite sides of the double bond.

As used throughout this application, the term "halogen" and the term "halo", unless otherwise stated, designate all four halogens, i.e. fluorine, chlorine, bromine and iodine. A preferred halogen is chlorine or fluorine. When R¹ and/or R² is halo, chlorine is especially preferred. When M is halo, fluorine is especially preferred.

As used herein, the term "lower alkyl" includes both straight chain and branched chain alkyl groups having from 1 to 7 carbon atoms, such as methyl, ethyl, propyl, isopropyl, preferably methyl and ethyl. With regard to R³, isopropyl and n-propyl are preferred, isobutyl is also preferred.

As used herein, the term "Halo lower alkyl" designates a lower alkyl group wherein one or more of the hydrogens is replaced by a halogen as defined above, which replacement can be at any site on the lower alkyl, including the end, such as chloroethyl. With regard to R³ fluoro lower alkyl is preferred.

As used herein, the term "Fluoro lower alkyl" designates a lower alkyl group wherein one or more of the hydrogens is replaced by fluorine, which replacement can be at any site on the lower alkyl, including the end, such as 1,1,1-trifluoroethane, 1,1,1-trifluoropropane and 1,1,1,3,3,3-hexafluoroisopropyl. A preferred fluoro lower alkyl group is 1,1,1,3,3,3-hexafluoroisopropyl.

The term "hydroxy lower alkyl" includes any hydroxy lower alkyl group where lower alkyl is defined as above. The hydroxy can be substituted at any place on the lower alkyl group such as hydroxy methyl, 1-hydroxy ethyl, 2-hydroxy propyl, 2-hydroxy isopropyl or 2-hydroxy-2-butyl. "Lower alkoxy lower alkyl" denotes any hydroxy lower alkyl group wherein the hydrogen of the hydroxy moiety is substituted by lower alkyl.

As used herein, "perfluoro-lower alkyl" means any lower alkyl group wherein all of the hydrogens of the lower alkyl group are substituted or replaced by fluoro. Among the preferred perfluoro-lower alkyl groups are trifluoromethyl, pentafluoroethyl, heptafluoropropyl, etc. An especially preferred perfluoro-lower alkyl group is trifluoromethyl.

As used herein, "lower alkyl thio" means a lower alkyl group as defined above where a thio group is bound to the rest of the molecule. Similarly "perfluoro-lower alkyl

thio” means a perfluoro-lower alkyl group as defined above where a thio group is bound to the rest of the molecule.

As used herein, “lower alkyl sulfonyl” means a lower alkyl group as defined above where a sulfonyl group is bound to the rest of the molecule, preferably lower alkyl

5 sulfonyl is methyl sulfonyl. Similarly “perfluoro-lower alkyl sulfonyl” means a perfluoro-lower alkyl group as defined above where a sulfonyl group is bound to the rest of the molecule

As used herein, “hydroxyamino” designates an amino group where one of the hydrogens is replaced by a hydroxy.

10 As used herein, “cycloalkyl” means a saturated hydrocarbon ring having from 3 to 8 carbon atoms, preferably from 5 to 6 carbon atoms, such as cyclopentyl and cyclohexyl. An especially preferred cycloalkyl is cyclopentyl.

As used herein, “heterocycloalkyl” means a saturated hydrocarbon ring having from 3 to 8 carbon atoms, preferably from 5 to 7 carbon atoms, and having one to two
15 heteroatoms which may be oxygen, sulfur or nitrogen. With regard to R^3 it is preferred to have a single heteroatom, preferably oxygen.

As used herein, “cycloalkenyl” means a cycloalkyl ring having from 3 to 8, and preferably from 5 to 7 carbon atoms, where one of the bonds between the ring carbons is unsaturated.

20 As used herein, the term “lower alkenyl” denotes an alkylene group having from 2 to 6 carbon atoms with a double bond located between any two adjacent carbons of the group, such as allyl and crotyl.

As used herein, the term “lower alkoxy” includes both straight chain and branched chain alkoxy groups having from 1 to 7 carbon atoms, such as methoxy, ethoxy, propoxy,
25 isopropoxy, preferably methoxy and ethoxy.

As used herein, the term “aryl” signifies aryl mononuclear aromatic hydrocarbon groups such as phenyl, tolyl, etc. which can be unsubstituted or substituted in one or more positions with halogen, nitro, lower alkyl, or lower alkoxy substituents and polynuclear aryl groups, such as naphthyl, anthryl, and phenanthryl, which can be unsubstituted or
30 substituted with one or more of the aforementioned groups. Preferred aryl groups are the substituted and unsubstituted mononuclear aryl groups, such as phenyl.

As used herein, the term "arylalkyl" denotes an alkyl group, preferably lower alkyl, in which one of the hydrogen atoms is replaced by an aryl group. Examples of arylalkyl groups are benzyl, 2-phenylethyl, 3-phenylpropyl, 4-chlorobenzyl, 4-methoxybenzyl and the like.

5 As used herein, the term "lower alkanolic acid" denotes lower alkanolic acids containing from 2 to 7 carbon atoms such as propionic acid, acetic acid and the like. The term "lower alkanoyl" denotes monovalent alkanoyl groups having from 2 to 7 carbon atoms such as propionoyl, acetyl and the like. The term "aroic acids" denotes aryl alkanolic acids where aryl is as defined above and alkanolic contains from 1 to 6 carbon atoms. The
10 term "aroyl" denotes aroic acids wherein aryl is as defined hereinbefore, with the hydrogen group of the COOH moiety removed. Among the preferred aroyl groups is benzoyl.

The heteroaromatic ring in R^4 can be an unsubstituted, mono- or di-substituted five- or six-membered heteroaromatic ring having from 1 to 3 heteroatoms selected from
15 the group consisting of oxygen, nitrogen or sulfur and connected by a ring carbon to the amine of the amide group shown. The heteroaromatic ring contains a first nitrogen heteroatom adjacent to the connecting ring carbon atom and if present, the other heteroatoms can be sulfur, oxygen or nitrogen. Heteroaromatic rings include, for example, pyrazinyl, pyridazinyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridinyl, pyrimidinyl,
20 thiadiazolyl (preferably 1,3,4-, 1,2,3-, 1,2,4-), triazinyl (preferably 1,3,5-, 1,2,4-), thiazolyl, oxazolyl, and imidazolyl. The preferred heteroaromatic rings which constitute R^4 are connected *via* a ring carbon atom to the amide group to form the amides of formula I.

R^4 is preferably an unsubstituted, mono- or di-substituted five- or six-membered,
25 heteroaromatic ring containing from 1 to 3 hetero atoms selected from the group consisting of nitrogen, oxygen or sulfur, with one hetero atom being nitrogen and connected to the remainder of the molecule by a ring carbon atom. In this case, the preferred rings are those which contain a nitrogen heteroatom adjacent to the connecting ring carbon. When R^4 is an unsubstituted, mono- or di-substituted five- or six-membered
30 heteroaromatic ring, the preferred rings are those which contain a nitrogen heteroatom adjacent to the connecting ring carbon and a second heteroatom adjacent to the connecting ring carbon or adjacent to said first heteroatom. Preferably R^4 is a five-membered

-10-

heteroaromatic ring. The preferred five-membered heteroaromatic rings contain 2 or 3 heteroatoms. Examples of such five-membered heteroaromatic rings are thiazolyl, imidazolyl, oxazolyl and thiadiazolyl, with thiazolyl being especially preferred.

When the heteroaromatic ring is a six-membered heteroaromatic, the ring is
5 connected by a ring carbon atom to the amine group shown, with one nitrogen heteroatom being adjacent to the connecting ring carbon atom. The preferred six-membered heteroaromatic rings include, for example, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, and triazinyl with pyridinyl being especially preferred.

Above heteroaromatic rings R^4 may optionally be mono- or di-substituted at a position on
10 a ring carbon atom other than adjacent to said connecting carbon atom with a substituent selected from the group consisting lower alkyl, halo, nitro, cyano, $-(CH_2)_n-OR^{20}$, $-(CH_2)_n-C(O)OR^{21}$, $-(CH_2)_n-CONHR^{22}$, $-(CH_2)_n-NHR^{23}$, with n , R^{20} , R^{21} , R^{22} and R^{23} being as defined above.

During the course of the synthetic sequence the various functional groups such as
15 the free carboxylic acid or hydroxy groups will be protected *via* conventional hydrolyzable ester or ether protecting groups. As used herein the term "hydrolyzable ester or ether protecting groups" designates any ester or ether conventionally used for protecting carboxylic acids or alcohols which can be hydrolyzed to yield the respective hydroxyl or carboxyl group. Exemplary ester groups useful for those purposes are those in which the
20 acyl moieties are derived from a lower alkanoic, aryl lower alkanoic, or lower alkane dicarboxylic acid. Among the activated acids which can be utilized to form such groups are acid anhydrides, acid halides, preferably acid chlorides or acid bromides derived from aryl or lower alkanoic acids. Example of anhydrides are anhydrides derived from monocarboxylic acid such as acetic anhydride, benzoic acid anhydride, and lower alkane
25 dicarboxylic acid anhydrides, e.g. succinic anhydride as well as chloro formats e.g. trichloro, ethylchloro formate being preferred. Suitable ether protecting groups for alcohols are, for example, the tetrahydropyranyl ethers such as 4-methoxy-5,6-dihydroxy-2H-pyran ethers. Others are aroylmethylethers such as benzyl, benzhydryl or trityl ethers or a-lower alkoxy lower alkyl ethers, for example, methoxymethyl or allylic ethers
30 or alkyl silylethers such as trimethylsilylether.


As used herein, the term "amino protecting group" designates any conventional amino protecting group which can be cleaved to yield the free amino group. The preferred

-11-


protecting groups are the conventional amino protecting groups utilized in peptide synthesis. Especially preferred are those amino protecting groups which are cleavable under mildly acidic conditions at about pH 3.0. Particularly preferred amino protecting groups are t-butoxycarbonyl (BOC), carbobenzyloxy (CBZ) and 9-

5 fluorenylmethoxycarbonyl(FMOC).

The compound of formula (I) of this invention constitutes two preferred species, i.e., the compound of formula



10 wherein A, B, R³ and R⁴ are as above;
and the compound of the formula



15 wherein A, M, R³ and R⁴ are as above.

In accordance with one preferable embodiment of the compound of formula I, R³ is lower alkyl having from 2 to 6 carbon atoms, preferred lower alkyl residues being isopropyl and n-propyl. In one preferred embodiment R³ is isopropyl, in another R³ is isobutyl. In another preferable embodiment R³ is -(CH₂)_s-V where "s" and "V" are as
20 defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms. In one preferred embodiment V is cyclopentyl, in another V is cyclohexyl. In another preferable embodiment R³ is halo lower alkyl having from 2 to 6 carbon atoms, preferred halo lower alkyl residues being fluoro lower alkyl as defined above.


25 Preferred R⁴ substituent in accordance with the present invention is where R⁴ is R¹⁷ as defined above. Further preferred R¹⁷ substituents are unsubstituted, mono- or di-

-12-

substituted thiazolyl, imidazolyl, oxazolyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, and triazinyl. In one preferred embodiment R^{17} is thiazolyl, in another R^{17} is pyridinyl. In accordance with further preferable embodiments, the heteroaromatic ring R^{17} is either unsubstituted, mono- or di-substituted independently with lower alkyl, halogen or $-(CH_2)_n-C(O)OR^{21}$, wherein n and R^{21} are as defined above. In another preferable embodiment M is hydrogen, halo or lower alkyl. Further preferred M substituents are hydrogen, fluoro, methyl or ethyl. In one preferred embodiment M is hydrogen, in another M is methyl.

In accordance with another preferable embodiment of the compound of formula

(I), A is



wherein Y and R^1 and R^2 are as defined above, the preferred Y substituent being sulfur. Preferred substituent R^1 is selected from the group consisting of hydrogen, halo, nitro, cyano, perfluoro lower alkyl, and lower alkyl sulfonyl. A further preferred R^1 substituent is selected from the group consisting of hydrogen, halo, nitro, cyano or perfluoro lower alkyl. In one preferred embodiment R^1 is halo, in another R^1 is hydrogen. Preferred substituent R^2 is selected from the group consisting of hydrogen, halo, nitro, cyano, perfluoro lower alkyl, lower alkyl sulfonyl, $R^{10}-[(CH_2)_y-W]_z$ and $R^{13}-(CH_2)_t-U$ where R^{10} , R^{13} , W , U , t , y and z are as defined above. Further preferred R^2 substituents are halo, lower alkyl sulfonyl, $R^{10}-[(CH_2)_y-W]_z$ and $R^{13}-(CH_2)_t-U$ where R^{10} , R^{13} , W , U , t , y and z are as defined above. In one preferred embodiment R^2 is lower alkyl sulfonyl, in another R^2 is $R^{10}-[(CH_2)_y-W]_z$ where R^{10} , W , U , y and z are as defined above, in another R^2 is $R^{13}-(CH_2)_t-U$ where R^{13} , U and t , are as defined above. In a further preferred embodiment R^2 is sulphonyl methyl, in another R^2 is $R^{10}-[(CH_2)_y-W]_z$ where W is SO_2 and R^{10} , U , y and z are as defined above.


In accordance with one embodiment of the compound of formula I-A, R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or

-13-

cyclohexyl (compound I-A1). Among the various embodiments of the compound I-A1 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen. Another embodiment of the compounds of formula (I-A) are those compounds where R³ is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-A2). Among the various embodiments of the compound I-A2 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-A, R⁴ is R¹⁷, which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R¹⁷ is a thiazole or pyridine ring (compound I-A3). Among the various embodiments of the compound I-A3 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A, A is



where Y is sulfur (compound I-A4). Among the various embodiments of the compound I-A4 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A, A is




where Y is sulfur (compound I-A5). Among the various embodiments of the compound I-A5 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

-14-


Among the embodiments of the compound of formula I-A1, R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A1(a)). Among the various embodiments of the compound I-A1(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A1, A is



where Y is sulfur (compound I-A1(b)). Among the various embodiments of the compound I-A1(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A1, A is




where Y is sulfur (compound I-A1(c)). Among the various embodiments of the compound I-A1(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A2, R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A2(a)). Among the various embodiments of the compound I-A2(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


-15-

Among the embodiments of the compound of formula I-A2, A is



- where Y is sulfur (compound I-A2(b)). Among the various embodiments of the compound I-A2(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A2, A is




- where Y is sulfur (compound I-A2(c)). Among the various embodiments of the compound I-A2(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

- Among the embodiments of the compound of formula I-A3, R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-A3(a)). Among the various embodiments of the compound I-A3(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

- Among the embodiments of the compound of formula I-A3 are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-A3(b)). Among the various embodiments of the compound I-A3(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A3, A is


-16-



where Y is sulfur (compound I-A3(c)). Among the various embodiments of the compound I-A3(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is

5 hydrogen.

Among the embodiments of the compound of formula I-A3, A is



where Y is sulfur (compound I-A3(d)). Among the various embodiments of the compound I-A3(d) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is

10 hydrogen.

Among the embodiments of the compound of formula I-A4, R³ can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl

15 (compound I-A4(a)). Among the various embodiments of the compound I-A4(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A4 are those compounds where R³ is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues

20 being isobutyl or isopropyl (compound I-A4(b)). Among the various embodiments of the compound I-A4(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A4, R⁴ is R¹⁷, which can

25 be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R¹⁷ is a thiazole or pyridine ring (compound I-A4(c)). Among the

-17-


various embodiments of the compound I-A4(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A5, R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-A5(a)). Among the various embodiments of the compound I-A5(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A5 are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-A5(b)). Among the various embodiments of the compound I-A5(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A5, R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A5(c)). Among the various embodiments of the compound I-A5(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-A1(a), A is



where Y is sulfur (compound I-A1(a-1)). Among the various embodiments of the compound I-A1(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A1(a), A is

-18-




where Y is sulfur (compound I-A1(a-2)). Among the various embodiments of the compound I-A1(a-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A1(b), R⁴ is R¹⁷, which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R¹⁷ is a thiazole or pyridine ring (compound I-A1(b-1)). Among the various embodiments of the compound I-A1(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A1(c), R⁴ is R¹⁷, which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R¹⁷ is a thiazole or pyridine ring (compound I-A1(c-1)). Among the various embodiments of the compound I-A1(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-A2(a), A is



where Y is sulfur (compound I-A2(a-1)). Among the various embodiments of the compound I-A2(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A2(a), A is

-19-




where Y is sulfur (compound I-A2(a-2)). Among the various embodiments of the compound I-A2(a-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A2(b), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A2(b-1)). Among the various embodiments of the compound I-A2(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A2(c), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A2(c-1)). Among the various embodiments of the compound I-A2(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-A3(a), A is



where Y is sulfur (compound I-A3(a-1)). Among the various embodiments of the compound I-A3(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-A3(a), A is

-20-




where Y is sulfur (compound I-A3(a-2)). Among the various embodiments of the compound I-A3(a-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A3(b), A is



where Y is sulfur (compound I-A3(b-1)). Among the various embodiments of the compound I-A3(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A3(b), A is



where Y is sulfur (compound I-A3(b-2)). Among the various embodiments of the compound I-A3(b-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A3 (c), R^3 can be $-(CH_2)_s-$ V where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-A3(c-1)). Among the various embodiments of the compound I-A3(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

-21-

Among the embodiments of the compound of formula I-A3(c) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-A3(c-2)). Among the various embodiments of the compound I-A3(c-2) are included those compounds where M is
5 hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A3 (d), R^3 can be $-(CH_2)_s-$ V where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl
10 (compound I-A3(d-1)). Among the various embodiments of the compound I-A3(d-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A3(d) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower
15 alkyl residues being isobutyl or isopropyl (compound I-A3(d-2)). Among the various embodiments of the compound I-A3(d-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A4(a), R^4 is R^{17} , which
20 can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A4(a-1)). Among the various embodiments of the compound I-A4(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more
25 particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A4(b), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those
30 compounds where R^{17} is a thiazole or pyridine ring (compound I-A4(b-1)). Among the various embodiments of the compound I-A4(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A4 (c), R^3 can be $-(CH_2)_s-$ V where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-A4(c-1)). Among the various embodiments of the compound I-A4(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A4(c) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-A4(c-2)). Among the various embodiments of the compound I-A4(c-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A5(a), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A5(a-1)). Among the various embodiments of the compound I-A5(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A5(b), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-A5(b-1)). Among the various embodiments of the compound I-A5(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-A5(c), R^3 can be $-(CH_2)_s-$ V where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-A5(c-1)). Among the various embodiments of the compound I-A5(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

-23-

Among the embodiments of the compound of formula I-A5(c) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-A5(c-2)). Among the various embodiments of the compound I-A5(c-2) are included those compounds where M is

5 hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

In accordance with one embodiment of the compound of formula I-B, R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or


10 cyclohexyl (compound I-B1). Among the various embodiments of the compound I-B1 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen. Another embodiment of the compounds of formula I-B are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or

15 isopropyl (compound I-B2). Among the various embodiments of the compound I-B2 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B, R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole,

20 pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-B3). Among the various embodiments of the compound I-B3 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


25 Among the embodiments of the compound of formula I-B, A is



where Y is sulfur (compound I-B4). Among the various embodiments of the compound I-B4 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

30 Among the embodiments of the compound of formula I-B, A is


-24-



where Y is sulfur (compound I-B5). Among the various embodiments of the compound I-B5 are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


- 5 Among the embodiments of the compound of formula I-B1, R⁴ is R¹⁷, which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R¹⁷ is a thiazole or pyridine ring (compound I-B1(a)). Among the various embodiments of the compound I-B1(a) are included those compounds where M is
- 10 hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B1, A is



- where Y is sulfur (compound I-B1(b)). Among the various embodiments of the compound
- 15 I-B1(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B1, A is




- 20 where Y is sulfur (compound I-B1(c)). Among the various embodiments of the compound I-B1(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

-25-


Among the embodiments of the compound of formula I-B2, R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-B2(a)). Among the various embodiments of the compound I-B2(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B2, A is



where Y is sulfur (compound I-B2(b)). Among the various embodiments of the compound I-B2(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B2, A is



where Y is sulfur (compound I-B2(c)). Among the various embodiments of the compound I-B2(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-B3, R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-B3(a)). Among the various embodiments of the compound I-B3(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B3 are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues

-26-


being isobutyl or isopropyl (compound I-B3(b)). Among the various embodiments of the compound I-B3(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

5 Among the embodiments of the compound of formula I-B3, A is



where Y is sulfur (compound I-B3(c)). Among the various embodiments of the compound I-B3(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is
 10 hydrogen.

Among the embodiments of the compound of formula I-B3, A is



where Y is sulfur (compound I-B3(d)). Among the various embodiments of the compound I-B3(d) are included those compounds where M is hydrogen, halo or lower alkyl,
 15 particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B4, R³ can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl
 20 (compound I-B4(a)). Among the various embodiments of the compound I-B4(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B4 are those compounds where R³ is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues
 25 being isobutyl or isopropyl (compound I-B4(b)). Among the various embodiments of the compound I-B4(b) are included those compounds where M is hydrogen, halo or lower

-27-

alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-B4, R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-B4(c)). Among the various embodiments of the compound I-B4(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B5, R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-B5(a)). Among the various embodiments of the compound I-B5(a) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B5 are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-B5(b)). Among the various embodiments of the compound I-B5(b) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B5, R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-B5(c)). Among the various embodiments of the compound I-B5(c) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


Among the embodiments of the compound of formula I-B1(a), A is



-28-

where Y is sulfur (compound I-B1(a-1)). Among the various embodiments of the compound I-B1(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

5 Among the embodiments of the compound of formula I-B1(a), A is



where Y is sulfur (compound I-B1(a-2)). Among the various embodiments of the compound I-B1(a-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is

10 hydrogen.


Among the embodiments of the compound of formula I-B1(b), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-B1(b-1)). Among the

15 various embodiments of the compound I-B1(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B1(c), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those

20 compounds where R^{17} is a thiazole or pyridine ring (compound I-B1(c-1)). Among the various embodiments of the compound I-B1(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


25 Among the embodiments of the compound of formula I-B2(a), A is



-29-

where Y is sulfur (compound I-B2(a-1)). Among the various embodiments of the compound I-B2(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

5 Among the embodiments of the compound of formula I-B2(a), A is



where Y is sulfur (compound I-B2(a-2)). Among the various embodiments of the compound I-B2(a-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


10 Among the embodiments of the compound of formula I-B2(b), R⁴ is R¹⁷, which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R¹⁷ is a thiazole or pyridine ring (compound I-B2(b-1)). Among the

15 various embodiments of the compound I-B2(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B2(c), R⁴ is R¹⁷, which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole,

20 pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R¹⁷ is a thiazole or pyridine ring (compound I-B2(c-1)). Among the various embodiments of the compound I-B2(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.


25 Among the embodiments of the compound of formula I-B3(a), A is



-30-


where Y is sulfur (compound I-B3(a-1)). Among the various embodiments of the compound I-B3(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

5 Among the embodiments of the compound of formula I-B3(a), A is




where Y is sulfur (compound I-B3(a-2)). Among the various embodiments of the compound I-B3(a-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is
10 hydrogen.

Among the embodiments of the compound of formula I-B3(b), A is



where Y is sulfur (compound I-B3(b-1)). Among the various embodiments of the compound I-B3(b-1) are included those compounds where M is hydrogen, halo or lower
15 alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B3(b), A is



where Y is sulfur (compound I-B3(b-2)). Among the various embodiments of the compound I-B3(b-2) are included those compounds where M is hydrogen, halo or lower
20 alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

Among the embodiments of the compound of formula I-B3 (c), R³ can be -(CH₂)_s-V where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl

-31-

group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-B3(c-1)). Among the various embodiments of the compound I-B3(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

5 Among the embodiments of the compound of formula I-B3(c) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-B3(c-2)). Among the various embodiments of the compound I-B3(c-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more
10 particularly where M is hydrogen.

 Among the embodiments of the compound of formula I-B3(d), R^3 can be $-(CH_2)_s-$ V where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-B3(d-1)). Among the various embodiments of the compound I-B3(d-1) are
15 included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

 Among the embodiments of the compound of formula I-B3(d) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-B3(d-2)). Among the various
20 embodiments of the compound I-B3(d-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

 Among the embodiments of the compound of formula I-B4(a), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole,
25 pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-B4(a-1)). Among the various embodiments of the compound I-B4(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

30 Among the embodiments of the compound of formula I-B4(b), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those

compounds where R^{17} is a thiazole or pyridine ring (compound I-B4(b-1)). Among the various embodiments of the compound I-B4(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

5 Among the embodiments of the compound of formula I-B4 (c), R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-B4(c-1)). Among the various embodiments of the compound I-B4(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where
10 M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

 Among the embodiments of the compound of formula I-B4(c) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-B4(c-2)). Among the various embodiments of the compound I-B4(c-2) are included those compounds where M is
15 hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

 Among the embodiments of the compound of formula I-B5(a), R^4 is R^{17} , which can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those
20 compounds where R^{17} is a thiazole or pyridine ring (compound I-B5(a-1)). Among the various embodiments of the compound I-B5(a-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

 Among the embodiments of the compound of formula I-B5(b), R^4 is R^{17} , which
25 can be an unsubstituted, mono- or di-substituted thiazole, imidazole, oxazole, thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, or triazine ring, and particularly those compounds where R^{17} is a thiazole or pyridine ring (compound I-B5(b-1)). Among the various embodiments of the compound I-B5(b-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more
30 particularly where M is hydrogen.

 Among the embodiments of the compound of formula I-B5(c), R^3 can be $-(CH_2)_s-V$ where "s" and "V" are as defined above, the preferred V substituent being a cycloalkyl


-33-

group which contains from 3 to 8 carbon atoms, preferably cyclopentyl or cyclohexyl (compound I-B5(c-1)). Among the various embodiments of the compound I-B5(c-1) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more particularly where M is hydrogen.

5 Among the embodiments of the compound of formula I-B5(c) are those compounds where R^3 is lower alkyl having from 2 to 6 carbon atoms, preferably lower alkyl residues being isobutyl or isopropyl (compound I-B5(c-2)). Among the various embodiments of the compound I-B5(c-2) are included those compounds where M is hydrogen, halo or lower alkyl, particularly where M is hydrogen, fluoro or methyl, more
10 particularly where M is hydrogen.

The compounds of the present invention may be prepared as is shown in the following reaction schemes.

In the case when Z equals



15 the preparation of the compounds is outlined in Scheme I. The corresponding alkyl iodides are reacted with a propargyl ester to give the substituted 2-iodo alkenoates (see Y. Ichinose et al., *Tet. Lett.* **1989**, 24, 3155). Alternatively, compounds where M equals hydrogen, can be prepared by reacting aldehydes R^3 -CHO with an appropriate Wittig reagent, preferably triethyl phosphono acetate, to give the acrylic esters, which then are
20 transformed into the 2-halo-alk-2-enoates by a halogenation / dehydro-halgenation procedure known in the art. Subsequent Pd(0)-catalyzed cross coupling with heteroaromatic boronic acids and boronic esters, respectively, yields the hetaryl-substituted alk-2-enoates of formula II. The latter intermediates can also be prepared from in situ formed olefinic boronic esters with the corresponding hetaryl bromides and iodides.


25 For the preparation of compounds of formula II where M equals halo, the (hetero)aryl substituted α -keto ester may be reacted with the appropriate halo-substituted organophosphonate using conditions known in the art. In order to prepare amides of formula III, these can be directly reacted with the deprotonated amines R^{17} -NH₂, using appropriate Grignard reagents, preferably isopropyl magnesium chloride. Alternatively,
30 alkenoates of formula II can be hydrolyzed to the corresponding acids of formula V by

-34-

known procedures of alkaline or acidic saponification. Subsequent activation of the acids by known means in the area of peptide coupling, followed by reaction with amines R^{17} - NH_2 also yield the amides of formula III. For the synthesis of amides of the formula VI, the acids are reacted with substituted ureas $H_2N-CO-NH-R^{16}$. Reduction of compounds of
5 formula III and VI can be carried out using hydride reagents, preferably sodium borohydride, to afford the saturated alkanoate amides of formula IV and VII.

-35-

Scheme I



-36-

Experimentals:**General Procedures:**

All water or air-sensitive reactions were conducted in dry solvents under an inert atmosphere. Mass spectra (MS) were obtained on a Agilent 1100 MSD spectrometer
 5 operating in electrospray mode or on a Agilent 5973N GC-MSD.

List of abbreviations:

	THF	tetrahydrofuran
	DMSO	dimethylsulfoxide
10	TBTU	O-benzotriazol-1-yl-N,N,N',N' tetramethyluroniumtetrafluoroborate
	HATU	O-(7-Aza-1-benzotriazolyl)-N,N,N',N'- tetramethyluroniumhexafluorophosphate
	DMF	N,N-dimethylformamide
	NMP	1-methyl-2-pyrrolidinone
15	brine	saturated aqueous sodium chloride solution
	r.t.	room temperature / ambient temperature
	TBME	2-methoxy-2-methyl-propane
	DMA	N,N-dimethyl-acetamide
	HPLC	high pressure liquid chromatography
20	TFA	trifluoroacetic acid
	NH ₃	ammonia
	HOAt	1-hydroxy-7-azabenzotriazole
	MTB ether	2-methoxy-2-methyl-propane

25 **General Procedure I for the preparation of substituted 2-iodo-alk-2-enoates:**

The alkyl iodides mentioned below were reacted with ethyl propiolate in the presence of triethylborane as described (Y. Ichinose et al., Tet. Lett. 1989, 30(24), 3155-3158)

(Z)-Ethyl 3-cyclohexyl-2-iodo-propenoate, MS(EI), M⁺ = 308,

30 reagent used: cyclohexyl iodide

(Z)-Ethyl 3-cyclopentyl-2-iodo-propenoate, MS(EI), M⁺ = 294,

reagent used: cyclopentyl iodide

-37-

(Z)-Ethyl 2-iodo-4-methyl-pent-2-enoate, MS(EI), $M^+ = 268$,
reagent used: 2-iodo-propane


General procedure II for the preparation of compounds of formula I:

- 5 Add the heteroaromatic boronic acid (1.2 equiv. to vinyl halide) to a stirred mixture of vinyl halide, palladium catalyst (5 mol%) and potassium carbonate (2.5 equiv) in toluene (3ml / 1.0 mmol vinyl halide) and heat the mixture at 70 °C for 14h. Subsequently filter the mixture, add water and extract with MTB ether. Combine the organic layers, dry over sodium sulfate and concentrate under reduced pressure. Dissolve the crude product in ethanol (2 ml/mmol), add 2N sodium hydroxide solution (2ml/mmol) and heat at 70 °C for 14h. Acidify with 1N hydrochloric acid, extract with MTB ether, dry. combined organic layers over sodium sulfate and concentrate under reduced pressure. Dissolve the crude product in THF (3ml/mmol), add TBTU (1.5 equiv), triethylamine (2 equiv.) and the respective amine (1.5 equiv.) and stir for 14h at ambient temperature.
- 10 Aqueous work-up using saturated sodium hydrogencarbonate solution / saturated citric acid and extraction with MTB ether gives the crude product. Purification by flash chromatography (silica) eluting with a mixture of hexanes / ethyl acetate yields the title compounds.

20 The following Examples illustrate compounds of the present invention and methods for their synthesis.

Example 1:

(E)-3-Cyclohexyl-2-(5-chloro-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide



- 25 a) (E)-3-Cyclohexyl-2-(5-chloro-thiophen-2-yl)-propenoic acid:
 reaction of 5-Chloro-thiophen-2-yl-boronic acid (0.63g, 3.9 mmol) with (Z)-Ethyl 3-cyclohexyl-2-iodo-propenoate (1.0g, 3.25 mmol) using Pd(dba)₂ as catalyst as outlined in

-38-


General Procedure II.

b) (E)-3-Cyclohexyl-2-(5-chloro-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide:
reaction of 2-aminothiazole (0.35, 3.47mmol) with compound 1a) according to General
Procedure II, 0.13g of a yellowish solid, $MS(M^+ + H) = 353$.

5

Example 2:


(E)-3-Cyclopentyl-2-(thiophen-3-yl)-N-thiazol-2'-yl-propenylamide



- a) (E)-3-Cyclopentyl-2-(thiophen-3-yl)-propenoic acid:
10 reaction of thiophen-3-yl boronic acid (0.16g, 1.22 mmol) with (Z)-Ethyl 3-cyclopentyl-2-
iodo-propenoate (0.30g, 1.02 mmol) using $Pd(dba)_2$ as catalyst as outlined in General
Procedure II.
- b) (E)-3-Cyclohexyl-2-(5-chloro-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide:
reaction of 2-aminothiazole (0.13g, 1.22 mmol) with compound 2a) according to General
15 Procedure II, 0.53g of a yellowish solid, $MS(M^+ + H) = 305$.

Example 3:


(E)-3-Cyclohexyl-2-(benzothiophen-2-yl)-N-thiazol-2'-yl-propenylamide



- 20 a) (E)-3-Cyclohexyl-2-(benzothiophen-2-yl)-propenoic acid :
reaction of thiophen-3-yl boronic acid (0.69 g, 3.90 mmol) with (Z)-Ethyl 3-cyclohexyl-2-
iodo-propenoate (1.0 g, 3.25 mmol) using $Pd(dba)_2$ as catalyst as outlined in General
Procedure II.
- b) (E)-3-Cyclohexyl-2-(benzothiophen-2-yl)-N-thiazol-2'-yl-propenylamide:
25 reaction of 2-aminothiazole (0.30g, 3.0 mmol) with compound 3a) according to General

-39-

Procedure II, 0.40g of a solid, $MS(M^+ + H) = 369$.


Example 4:**(E)-3-Cyclopentyl-2-(benzothiophen-2-yl)-N-thiazol-2'-yl-propenylamide**

5

a) (E)-3-Cyclopentyl-2-(benzothiophen-2-yl)-propenoic acid

reaction of benzothiophen-2-yl boronic acid (0.22g, 1.22 mmol) with (Z)-Ethyl 3-cyclopentyl-2-iodo-propenoate (0.30g, 1.02 mmol) using $Pd(dba)_2$ as catalyst as outlined in General Procedure II.

- 10 b) (E)-3-Cyclopentyl-2-(benzothiophen-2-yl)-N-thiazol-2'-yl-propenylamide:
reaction of 2-aminothiazole (0.11g, 1.06 mmol) with compound 4a) according to General Procedure II, 0.05g of a yellowish solid, $MS(M^+ + H) = 355$.

Example 5:15 **(E)-3-Cyclohexyl-2-(benzofuran-2-yl)-N-thiazol-2'-yl-propenylamide**

a) (E)-3-Cyclohexyl-2-(benzofuran-2-yl)-propenoic acid:


- 20 reaction of benzofuran-2-yl-boronic acid (0.63g, 3.89 mmol) with (Z)-Ethyl 3-cyclohexyl-2-iodo-propenoate (1.0g, 3.25 mmol) using $Pd(dba)_2$ as catalyst as outlined in General Procedure II.

b) (E)-3-Cyclohexyl-2-(benzofuran-2-yl)-N-thiazol-2'-yl-propenylamide:

reaction of 2-aminothiazole (0.28, 2.80 mmol) with compound 5a) according to General Procedure II, 0.18 g of a yellowish solid, $MS(M^+ + H) = 353$.

25

-40-

Example 6:**(E)-4-Methyl-2-(5-bromo-thiophen-2-yl)-N-thiazol-2'-yl-pent-2-enylamide.**


- 5 a) (E)-4-Methyl-2-(5-bromo-thiophen-2-yl)-propenoic acid:

5-Bromo-thiophen-2-yl-zinc bromide (7.46 ml of a 0.5M solution in THF, 3.73 mmol) was added under argon atmosphere to a mixture containing (Z)-Ethyl 2-iodo-4-methyl-pent-2-enoate (0.5 g, 1.86 mmol), Pd(dppf)Cl₂ (136 mg, 10 mol%) and copper(I) iodide (30mg, 10 mol%) in 5ml of anhydr. THF. The mixture was stirred at reflux temperature for 14h.

- 10 After cooling, the reaction was quenched with 1N hydrochloric acid, followed by extraction with methylene dichloride. Hydrolysis of the ester was achieved as described according to the General procedure II.


- b) (E)-3-Cyclohexyl-2-(5-chloro-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide: reaction of 2-aminothiazole (0.16, 1.60 mmol) with compound 8a) according to General

- 15 Procedure II, 0.02 g of a yellowish solid, MS(M⁺+H)=358.


Example 7:**(E)-3-Cyclohexyl-2-(5-chloro-thiophen-2-yl)-N-thiazol-2'-yl-propionamide**

- 20 Compound 1b (0.11 g, 0.31 mmol) was dissolved in 2 ml of dry THF/EtOH (1:1), sodium borohydride (0.05 g, 1.32 mmol) was added at ambient temperature, and the reaction mixture was stirred for 14h. Aqueous work-up and extraction with ethyl acetate gave a yellowish gum after evaporation, which crystallized from acetonitrile to yield 22 mg of a colorless solid, MS(M⁺+H)=355.

-41-

Example 8:**(E)-3-Cyclohexyl-2-(benzothiophen-2-yl)-N-thiazol-2'-yl-propionamide**

Compound 3b) (0.23 g, 0.62 mmol) was dissolved in 4 ml of dry THF/EtOH (1:1) and sodium borohydride (0.095 g, 2.51 mmol) was added at ambient temperature and stirred for 14h. Aqueous work-up and extraction with ethyl acetate gave a yellowish gum after evaporation, which crystallized from acetonitrile to yield 85 mg of a colorless solid, MS(M⁺+H)=371.

Example 9:**(E)-3-Cyclopentyl-2-(5-methylsulfonyl-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide**


- a) Ethyl (E)-3-Cyclopentyl-2-(5-methylsulfonyl-thiophen-2-yl)-propenoate:
Ethyl (Z)-3-cyclopentyl-2-iodo-propenoate (0.1 g, 0.34 mmol) was dissolved in 2 ml of DMSO. Bis(pinacolato)diborane (0.093 g, 0.37 mmol), potassium acetate (0.1 g, 1.02 mmol) and Pd(dppf)Cl₂ (0.02g) were added at ambient temperature and then stirred for 2h at 80°C under argon atmosphere. After cooling, the mixture was added dropwise via a syringe to a solution of 2-iodo-5-methylsulfonyl-thiophene (0.036g, 0.12 mmol) in 2M sodium carbonate (0.29 ml) / DMSO (1 ml) at 50°C and stirred for an additional 1.5h.
- 20 Aqueous work-up, extraction with ethyl acetate, drying over sodium sulfate and evaporation gave the crude product.
- b) (E)-3-Cyclopentyl-2-(5-methylsulfonyl-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide:

-42-

To a solution of 2-aminothiazole (0.048g, 0.48 mmol) in THF (1 ml) was added isopropylmagnesium chloride (0.24 ml of a 2M solution in THF, 0.48 mmol) at -35°. The mixture was warmed and stirred under argon at ambient temperature for 1h. After cooling to -35°C, crude compound 9a) (0.09g, 0.12 mmol) was added dropwise and stirred at 50°C overnight. Work-up with sat. ammonium chloride solution, extraction with ethyl acetate, drying over sodium sulfate and evaporation gave the crude product. RP-chromatography using water/acetonitrile as eluent system gave 15 mg of a yellowish solid, MS(M⁺+H)=383.

10 Example 10:

(E)-3-Cyclopentyl-2-(5-sulfamoyl-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide



- a) 5-Bromo-thiophen-2-sulfonic acid dimethylaminomethylene amide: to a solution of 5-bromo-2-thiophen sulfonamide (0.3 g, 1.24 mmol) in abs. DMF (3 ml) was added N,N-dimethylformamide dimethylacetal (0.17 ml, 1.24 mmol), and the reaction mixture was stirred at ambient temperature overnight. Aqueous work-up, extraction with ethyl acetate, drying over sodium sulfate and evaporation gave the crude product as a colorless crystalline solid.
- b) Ethyl (E)-3-Cyclopentyl-2-[(5-dimethylaminomethylene sulfamoyl)-thiophen-2-yl]-propenoate: processed as described for compound 9a), using (Z)-Ethyl 3-cyclopentyl-2-iodo-propenoate (0.44g, 1.49 mmol), compound 12a) (0.35g), bis(pinacolato)diborane (0.42 g, 1.64 mmol), potassium acetate (0.44 g, 4.47 mmol) and Pd(dppf)Cl₂ (0.066g).
- c) (E)-3-Cyclopentyl-2-[(5-dimethylaminomethylene sulfamoyl)-thiophen-2-yl]-N-thiazol-2'-yl-propenoate: processed as described for compound 9b), using compound 10b) (0.61g, 1.59 mmol), 2-aminothiazole (0.47g, 4.72 mmol), isopropylmagnesium chloride (2.36 ml of a 2M solution in THF, 4.72 mmol).
- d) (E)-3-Cyclopentyl-2-(5-sulfamoyl-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide: Compound 10c) (0.04g, 0.09mmol) was dissolved in a solution of ethanol / 5N HCl aq (5


-43-

ml, 1:1 v:v) and heated at 80°C for 2h. Subsequent work-up with sat. sodium hydrogencarbonate solution and extraction with ethyl acetate yielded the crude product. Flash chromatography using methylene chloride/ethanol 98/2 as eluent system gave the product as a colorless solid, MS(M⁺+H)=384.

5

Example 11:

(E)-3-Cyclopentyl-2-(5-N-morpholinosulfonyl-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide




- 10 a) Ethyl (E)-3-Cyclopentyl-2-(5-N-morpholinosulfonyl-thiophen-2-yl)-propenoate: processed as described for compound 9a), using (Z)-Ethyl 3-cyclopentyl-2-iodo-propenoate (0.1g, 0.36 mmol), 2-bromo-5-N-morpholinosulfonyl-thiophen (0.078g, 0.25 mmol), bis(pinacolato)diborane (0.42 g, 1.64 mmol), potassium acetate (0.11 g, 1.1 mmol) and Pd(dppf)Cl₂ (0.04g).
- 15 b) (E)-3-Cyclopentyl-2-(5-N-morpholinosulfonyl-thiophen-2-yl)-N-thiazol-2'-yl-propenylamide: processed as described for compound 9a), using compound 11a) (0.2g, 0.25 mmol), 2-aminothiazole (0.14g, 1.44 mmol), isopropylmagnesium chloride (0.72 ml of a 2M solution in THF, 1.44 mmol). Flash chromatography using hexanes/ethyl ester 9/1 as eluent system gave the product, MS(M⁺+H)=454.

20

Example 12:

(Z)-3-Cyclopentyl-N-thiazol-2-yl-2-thiophen-3-yl-acrylamide:



-44-

- a) (Z)-3-Cyclopentyl-2-thiophen-3-yl-acrylic acid: 5-Bromo-thiophen-2-yl-zinc bromide (20.4 ml of a 0.5M solution in THF, 10.2 mmol) was added under argon atmosphere to a mixture containing (Z)-ethyl 2-iodo-4-methyl-pent-2-enoate (1.5 g, 5.1 mmol), Pd(dppf)Cl₂ (373 mg, 10 mol%) and copper(I) iodide (60 mg, 16 mol%) in 5 ml of anhydr. THF. The mixture was stirred at 75°C for 14h. After cooling, the reaction mixture was quenched with 1N hydrochloric acid, and extracted with MTB ether to afford 1.5 g of a red residue after removal of volatiles. This material was taken up in a mixture of EtOH (20 mL) and 2 N aqueous sodium hydroxide solution (20 mL). The resulting reaction mixture was stirred at 75°C for 14h, cooled, and poured onto 1 N aqueous hydrochloric acid. The resulting mixture was extracted with MTB ether. The extracts were dried over sodium sulfate and concentrated to afford 1.24 g of impure (Z)-3-cyclopentyl-2-thiophen-3-yl-acrylic acid.
- b) (Z)-3-Cyclopentyl-N-thiazol-2-yl-2-thiophen-3-yl-acrylamide: reaction of 2-aminothiazole (0.16 g, 1.62 mmol) with compound 12a) (0.3 g, 1.35 mmol) according to General Procedure II, 94 mg of a solid, MS(M⁺+H)=305

The compounds of the present invention may be used as medicaments in human or veterinary medicine. The compounds may be administered by various routes, for example, by oral or rectal routes, topically or parenterally, for example by injection, and are usually employed in the form of a pharmaceutical composition.

Such compositions may be prepared by methods well known in the pharmaceutical art and normally comprise at least one active compound in association with a pharmaceutically acceptable diluent or carrier. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier or diluted by a carrier, and/or enclosed within a carrier which may, for example, be in the form of a capsule, sachet, paper or other container. Where the carrier serves as a diluent, it may be solid, semi-solid, or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition may be in the form of tablets, lozenges, sachets, cachets, elixirs, suspensions, solutions, syrups, aerosol (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, injection solutions and suspensions and sterile packaged powders.

-45-

Some examples of suitable carriers are lactose, dextrose, vegetable oils, benzyl alcohols, alkylene glycols, polyethylene glycols, glycerol triacetate, gelatin, carbohydrates such as starch and petroleum jelly, sucrose sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, syrup, methyl cellulose, methyl- and propyl- hydrobenzoate, talc, magnesium stearate and mineral oil. The compounds of formula (I) can also be lyophilized and the lyophilizates obtained used, for example, for the production of injection preparations. The preparations indicated can be sterilized and/or can contain auxiliaries such as lubricants, preservatives, stabilizers and/or wetting agents, emulsifiers, salts for affecting the osmotic pressure, buffer substances, colourants, flavourings and/or one or more further active compounds, e.g. one or more vitamins. Compositions of the invention may be formulated so as to provide, quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 to about 500 mg, more usually about 25 to about 300 mg, of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary doses for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier.

Example A

Tablets containing the following ingredients can be produced in a conventional manner:

Ingredients (mg per capsule)

Compound of formula I	10.0 - 100.0
Lactose	125.0
Corn starch	75.0
Talc	4.0
Magnesium stearate	1.0

Example B

Capsules containing the following ingredients can be produced in a conventional manner:

-46-

Ingredients (mg per capsule)

Compound of formula I	25.0
Lactose	150.0
Corn starch	20.0
5 Talc	5.0

The pharmacological profile of the present compounds may be demonstrated as follows:

An enzymatic coupled glucokinase (GK) assay using purified recombinant human islet GK was used to evaluate effects of the activators. In this assay, GK catalyzes glucose phosphorylation in the presence of ATP. The product of this reaction, glucose-6-phosphate, is then oxidized by an excess of glucose-6-dehydrogenase to produce gluconate-6-phosphate with concomitant reduction of NAD^+ to NADH (Davidson and Arion, 1987). The following outlines the two reactions involved:

Glucose + ATP \rightarrow Glucose-6-P + ADP (Glucokinase)

Glucose-6-P + NAD \rightarrow Gluconate-6-P + NADH (glucose-6-P dehydrogenase)

The NADH production detected by absorbance at 340nm is used to monitor the enzymatic activity.

The human islet GK isoform was expressed in *E.coli* as (His)₆-tagged fusion protein and purified with metal chelate affinity chromatography (Tiedge et al., 1997). After purification the enzyme was stored in aliquots at concentration 0.8 mg/ml in 25 mM NaH_2PO_4 , 150 mM NaCl, 100 mM imidazole, 1 mM DTT, 50 % glycerol at -80°C.

The assay was performed in flat bottom 96-well plates in a final incubation volume of 100 μ l. The incubation mixture consisted of 25 mM HEPES (pH7.4), 50 mM KCl, 2.5 mM MgCl_2 , 2 mM dithiothreitol, 4 U/ml glucose-6-phosphate dehydrogenase from *Leuconostoc mesenteroides*, 5 mM ATP, 1 mM NAD and 10 mM glucose. All reagents were from Sigma-Aldrich Co. (St. Louis, MO). Test compounds were dissolved in DMSO and then added to the reaction mixture giving the final DMSO concentration of 10%.

-47-

The reaction was initiated by addition of 20 μ l GK and run for 20 min at 37°C. The amount of formed NADH was measured as an increase in absorbance at 340 nm using a microplate reader.

The concentration of activator that produced 50% of the maximum increase in the activity of GK (EC_{50}) was calculated. The preferred compounds of formula I described within the examples have an EC_{50} less than or equal to 30 μ M.

EXAMPLE	EC_{50} (μ M)
1	1.840
3	1.532
5	1.824
9	2.002
10	1.263
11	0.909
12	5.598

EC_{50} values shown in the above table are at 10mM glucose.

10

References:


- Davidson A.L. and Arion W.J. Factors underlying significant underestimations of glucokinase activity in crude liver extracts: physiological implications of higher cellular activity. Arch. Biochem. Biophys. 253, 156-167, 1987.
- 15 Tidge M, Krug U. and Lenzen S. Modulation of human glucokinase intrinsic activity by SH reagents mirrors post-translational regulation of enzyme activity. Biochem. Biophys. Acta 1337, 175-190, 1997.

20


-48-

Claims


1. A compound of formula



wherein Z is



- 10 R^3 is lower alkyl or halo lower alkyl having from 2 to 6 carbon atoms or arylalkyl or $-(\text{CH}_2)_s\text{-V}$ where V is a 3 to 8-membered ring which is cycloalkyl, cycloalkenyl, or heterocycloalkyl having one heteroatom selected from oxygen and sulfur;
 s is independently 0, 1 or 2;
 M is hydrogen or halo or lower alkyl or perfluoro-lower alkyl;
- 15 A is



where Y is oxygen or sulfur, and

- R^1 and R^2 are independently hydrogen, halo, amino, hydroxyamino, nitro, cyano,
 20 sulfonamido, lower alkyl, $-\text{OR}^5$, $-\text{COOR}^5$, perfluoro-lower alkyl, lower alkyl thio,
 perfluoro-lower alkyl thio, lower alkyl sulfonyl, perfluoro-lower alkyl sulfonyl, lower
 alkyl sulfinyl,
 R^5 is hydrogen, lower alkyl or perfluoro-lower alkyl; or furthermore
 R^2 can be $-(\text{CH}_2)_n\text{-NR}^6\text{R}^7$, with $n=1,2,3$ or 4, and

-49-

- R^6 and R^7 are independently hydrogen or lower alkyl; or together with the nitrogen atom to which they are attached form a five or six-membered heteroaromatic ring containing from 1 to 3 heteroatoms selected from sulfur, oxygen or nitrogen; or a saturated 5- or 6-membered cycloheteroalkyl ring, which contains from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen; or
- R^2 can be alkynyl, substituted with hydrogen, lower alkyl, hydroxy lower alkyl, lower alkoxy lower alkyl, an unsubstituted or hydroxy substituted cycloalkyl ring containing 5 or 6 carbon atoms, a five- or six-membered saturated heterocyclic ring which contains from 1 to 3 hetero atoms selected from the group consisting of sulfur, oxygen or nitrogen, or an unsubstituted five- or six-membered heteroaromatic ring, connected by a ring carbon atom, which contains from 1 to 3 heteroatoms in the ring selected from the group consisting of sulfur, nitrogen and oxygen, or $-(CH_2)_n-NR^8R^9$, with $n=1,2$, and
- R^8 and R^9 are independently hydrogen or lower alkyl; or together with the nitrogen atom to which they are attached form a five or six-membered heteroaromatic ring containing from 1 to 3 heteroatoms selected from sulfur, oxygen or nitrogen; or a saturated 5- or 6-membered cycloheteroalkyl ring, which contains from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen; or
- R^2 can be $R^{10}-[(CH_2)_y-W]_z-$, with
- W is oxygen, sulfur, $-SO-$, $-SO_2-$, and
- R^{10} is a heteroaromatic ring, connected by a ring carbon atom, which contains from 5 to 6 ring members with from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur or nitrogen, or
- aryl containing 6 or 10 ring carbon atoms, or
- aryl containing 6 ring carbon atoms fused with a heteroaromatic ring containing 5 or 6 ring members with 1 or 2 heteroatoms in the ring being selected from the group consisting of nitrogen, oxygen or sulfur, or
- a saturated 5- or 6-membered cycloheteroalkyl ring, which contains from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen, or
- a cycloalkyl ring having 5 or 6 carbon atoms, or
- $-NR^{11}R^{12}$, with R^{11} and R^{12} being independently hydrogen or lower alkyl; y is independently 0,1,2,3 or 4; z is independently 0,1; or

-50-

- R^2 can be $R^{13}-(CH_2)_t-U-$, with
U is $-NHCO-$, $-CONH-$, $-NHSO_2-$, $-SO_2NH-$ and
 R^{13} in the same meaning of R^{10} and
perfluoro-lower alkyl, lower alkyl, lower alkoxy carbonyl or
- 5 $-NR^{14}R^{15}$, R^{14} and R^{15} are independently hydrogen or lower alkyl; or together with the
nitrogen atom to which they are attached form a five or six-membered heteroaromatic ring
containing from 1 to 3 heteroatoms selected from sulfur, oxygen or nitrogen; or a saturated
5- or 6- membered heterocycloalkyl ring, which contains from 1 to 2 heteroatoms selected
from the group consisting of oxygen, sulfur and nitrogen;
- 10 t is an integer being 0, 1, 2, 3 or 4;
 R^4 is $-C(O)NHR^{16}$, or is R^{17} ;
 R^{16} is hydrogen, lower alkyl, lower alkenyl, hydroxy lower alkyl,
 $-(CH_2)_n-COOR^{18}$, $-CO-(CH_2)_n-COOR^{19}$;
 R^{17} is an unsubstituted, mono- or di-substituted five- or six-membered heteroaromatic ring
- 15 connected by a ring carbon atom to the amide group shown, which five- or six-membered
heteroaromatic ring contains from 1 to 4 heteroatoms selected from sulfur, oxygen or
nitrogen, with one heteroatom being nitrogen which is adjacent to the connecting ring
carbon atom; said mono- or di-substituted heteroaromatic ring being mono- or di-
substituted at a position on a ring carbon atom other than adjacent to said connecting
- 20 carbon atom with a substituent selected from the group consisting of lower alkyl, halo,
nitro, cyano, $-(CH_2)_n-OR^{20}$, $-(CH_2)_n-COOR^{21}$,
 $-(CH_2)_n-CONHR^{22}$, $-(CH_2)_n-NHR^{23}$,
n is 0, 1, 2, 3 or 4;
 R^{18} , R^{19} , R^{20} , R^{21} , R^{22} and R^{23} are independently hydrogen or lower alkyl,
- 25 and its pharmaceutically acceptable salts thereof.

2. A compound according to claim 1, wherein
 R^4 is an unsubstituted, mono- or di-substituted five- or six-membered heteroaromatic ring
connected by a ring carbon atom to the amide group shown, which five- or six-membered
- 30 heteroaromatic ring contains from 1 to 4 heteroatoms selected from sulfur, oxygen or
nitrogen, with one heteroatom being nitrogen which is adjacent to the connecting ring
carbon atom; said mono- or di-substituted heteroaromatic ring being mono- or di-

-51-

substituted at a position on a ring carbon atom other than adjacent to said connecting carbon atom with a substituent selected from the group consisting of lower alkyl, halo, nitro, cyano, $-(CH_2)_n-OR^{20}$, $-(CH_2)_n-COOR^{21}$, $-(CH_2)_n-CONHR^{22}$, $-(CH_2)_n-NHR^{23}$,

5 n is 0, 1, 2, 3 or 4;

R^{20} , R^{21} , R^{22} and R^{23} are independently hydrogen or lower alkyl, and its pharmaceutically acceptable salts thereof.

3. A compound according to any of claims 1 to 2, wherein R^4 is an unsubstituted,
10 mono- or di-substituted five- or six-membered heteroaromatic ring selected from the group consisting of thiazolyl, imidazolyl, oxazolyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, or triazinyl.

15 4. A compound according to any of claims 1 to 3 wherein R^4 is thiazolyl or pyridinyl, unsubstituted, mono- or di-substituted independently by halogen, lower alkyl or $(CH_2)_n-C(O)OR^{21}$, wherein n is 0, 1 or 2 and R^{21} is lower alkyl.

5. A compound according to claim 1, wherein
20 R^4 is $-C(O)NHR^{16}$;
 R^{16} is hydrogen, lower alkyl, lower alkenyl, hydroxy lower alkyl, $-(CH_2)_n-COOR^{18}$, $-CO-(CH_2)_n-COOR^{19}$;
n is 0, 1, 2, 3 or 4;
 R^{18} and R^{19} are independently hydrogen or lower alkyl,
25 and its pharmaceutically acceptable salts thereof.

6. A compound according to any of claims 1 or 5, wherein R^4 is $-C(O)NHR^{16}$, and R^{16} is lower alkyl or lower alkenyl.

30 7. A compound according to any of claims 1 to 6, wherein R^1 is hydrogen, halo, nitro or cyano.

-52-

8. A compound according to any of claims 1 to 7, wherein R^1 is hydrogen or halo.
9. A compound according to any of claims 1 to 8, wherein R^2 is hydrogen, halo, nitro, cyano, sulfonamido, lower alkyl, $-OR^5$, $-COOR^5$, perfluoro-lower alkyl, lower alkyl
- 5 sulfonyl; or
- R^2 can be $R^{10}-[(CH_2)_y-W]_z-$, where
- W is oxygen, sulfur, $-SO-$, or $-SO_2-$, and
- R^{10} is a heteroaromatic ring, connected by a ring carbon atom, which contains from 5 to 6 ring members with from 1 to 2 heteroatoms selected from the group consisting of oxygen,
- 10 sulfur or nitrogen, or
- aryl containing 6 or 10 ring carbon atoms, or
- aryl containing 6 ring carbon atoms fused with a heteroaromatic ring containing 5 or 6 ring members with 1 or 2 heteroatoms in the ring being selected from the group consisting of nitrogen, oxygen or sulfur, or
- 15 a saturated 5- or 6-membered cycloheteroalkyl ring, which contains from 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen, or
- a cycloalkyl ring having 5 or 6 carbon atoms, or
- $-NR^{11}R^{12}$, with R^{11} and R^{12} being independently hydrogen or lower alkyl;
- y is independently 0,1,2,3 or 4; z is independently 0,1; or
- 20 R^2 can be $R^{13}-(CH_2)_t-U-$, with
- U is $-NHCO-$, $-CONH-$, $-NHSO_2-$, $-SO_2NH-$ and
- R^{13} in the same meaning of R^{10} and
- perfluoro-lower alkyl, lower alkyl, lower alkoxy carbonyl or
- $-NR^{14}R^{15}$, R^{14} and R^{15} are independently hydrogen or lower alkyl; or together with the
- 25 nitrogen atom to which they are attached form a five or six-membered heteroaromatic ring containing from 1 to 3 heteroatoms selected from sulfur, oxygen or nitrogen;
- t is an integer from 0 to 4.
10. A compound according to any of claims 1 to 9, wherein R^2 is halo, lower alkyl
- 30 sulfonyl or $R^{10}-[(CH_2)_y-W]_z-$.

-53-

11. A compound according to any of claims 1 to 10, wherein R^2 is sulfonylmethyl or $R^{10}-[(CH_2)_y-W]_z$ - where W is SO_2 .


12. A compound according to any of claims 1 to 11, wherein V is cyclopentyl,
5 cyclohexyl or cycloheptyl.

13. A compound according to any of claims 1 to 12, wherein V is cyclopentyl.

14. A compound according to any of claims 1 to 11, wherein R^3 is isopropyl or n-
10 propyl.


15. A compound according to any of claims 1 to 11, wherein R^3 is isobutyl.

16. A compound according to any of claims 1 to 15, wherein Z is



17. A compound according to claim 16, wherein M is hydrogen, fluorine or methyl.

18. A compound according to any of claims 1 to 17, wherein A is



19. A compound according to any of claims 1 to 18, wherein Y is sulfur.

20. The use of the compounds according to any of claims 1 to 19, or a
25 pharmaceutically acceptable salt thereof, for the treatment or prophylaxis of type II
diabetes.

-54-

21. A pharmaceutical composition comprising a compound of any of claims 1 to 19, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable diluent or carrier.

5

22. The use of a compound according to any of claims 1 to 19, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment or prophylaxis of type II diabetes.

10 23. A method for the prophylactic or therapeutic treatment of type II diabetes, which comprises administering a compound of any of claims 1 to 19, or a pharmaceutically acceptable salt thereof, to a human being or animal in need thereof.

15 24. A pharmaceutical composition for treating type II diabetes containing as an active ingredient a compound of any of claims 1 to 19, or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 03/37089

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D417/12 A61K31/427 A61P3/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01/12189 A (BRASCA MARIA GRABRIELLA ; TRAQUANDI GABRIELLA (IT); ORSINI PAOLO (IT);) 22 February 2001 (2001-02-22) claim 1	1
Y	WO 01/44216 A (HOFFMANN LA ROCHE) 21 June 2001 (2001-06-21) * Examples, claim 1 *	1-24
P,Y	EP 1 336 607 A (NOVO NORDISK AS) 20 August 2003 (2003-08-20) *Examples, claim 1*	1-24
Y	WO 02/14312 A (HOFFMANN LA ROCHE) 21 February 2002 (2002-02-21) *Examples, claim 1*	1-24
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

30 March 2004

Date of mailing of the international search report

15/04/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Baston, E

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 03/37089

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 02/08209 A (HOFFMANN LA ROCHE) 31 January 2002 (2002-01-31) *Examples, claim1* -----	1-24
Y	WO 01/85707 A (HOFFMANN LA ROCHE) 15 November 2001 (2001-11-15) *Examples, claim 1* -----	1-24
Y	WO 02/48106 A (HOFFMANN LA ROCHE) 20 June 2002 (2002-06-20) *Examples, claim1* -----	1-24
P,Y	WO 03/055482 A (NOVO NORDISK AS) 10 July 2003 (2003-07-10) claim 1 -----	1-24
Y	WERMUTH ET AL: "The Practise of Medicinal Chemistry" PRACTICE OF MEDICINAL CHEMISTRY, XX, XX, 1996, pages 203-237, XP002190259 table 13.2 -----	1-24
X	YOSHINA, S.; TANAKA,A.; USUI, T.: "Studies on heterocyclic compounds. XXIX. Synthesis of 2-'alpha-(N-substituted carbamoyl)-beta-(5-nitro-2-furyl)vinyl!fur ans and 2-'alpha-(N-substituted ureido)-beta-(5-nitro-2-furyl)vinyl! furans" YAKUGAKU ZASSHI, vol. 97, no. 9, 1977, pages 1007-1012, XP0009028638 examples III-11,III-15,III-16; table I -----	1

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 03/37089

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0112189 A	22-02-2001	AU 4971400 A	13-03-2001
		BG 106480 A	30-09-2002
		BR 0013143 A	11-06-2002
		CA 2383555 A1	22-02-2001
		CN 1373662 T	09-10-2002
		CZ 20020422 A3	13-11-2002
		EE 200200065 A	15-04-2003
		EP 1202733 A1	08-05-2002
		HR 20020128 A1	30-04-2003
		HU 0203542 A2	28-05-2003
		JP 2003507329 T	25-02-2003
		NO 20020684 A	03-04-2002
		PL 353455 A1	17-11-2003
		SK 1812002 A3	04-02-2003
		WO 0112189 A1	22-02-2001
		US 6218418 B1	17-04-2001
		ZA 200201511 A	11-03-2003
WO 0144216 A	21-06-2001	AU 2365201 A	25-06-2001
		CA 2392903 A1	21-06-2001
		CN 1411453 T	16-04-2003
		CZ 20022412 A3	16-10-2002
		WO 0144216 A1	21-06-2001
		EP 1242397 A1	25-09-2002
		HU 0203753 A2	28-03-2003
		JP 2003516980 T	20-05-2003
		NO 20022863 A	14-06-2002
		US 6353111 B1	05-03-2002
EP 1336607 A	20-08-2003	EP 1336607 A1	20-08-2003
		WO 03047626 A1	12-06-2003
		WO 03055482 A1	10-07-2003
		US 2003138416 A1	24-07-2003
		US 2003171411 A1	11-09-2003
WO 0214312 A	21-02-2002	AU 8399801 A	25-02-2002
		BR 0113312 A	01-07-2003
		CA 2419416 A1	21-02-2002
		CN 1466584 T	07-01-2004
		WO 0214312 A1	21-02-2002
		EP 1311504 A1	21-05-2003
		US 2002065275 A1	30-05-2002
		US 2002035266 A1	21-03-2002
		US 2002035267 A1	21-03-2002
WO 0208209 A	31-01-2002	AU 8760001 A	05-02-2002
		BR 0112658 A	24-06-2003
		CA 2416229 A1	31-01-2002
		CN 1443177 T	17-09-2003
		WO 0208209 A1	31-01-2002
		EP 1305301 A1	02-05-2003
		JP 2004504388 T	12-02-2004
		US 2002198200 A1	26-12-2002
		US 2002042512 A1	11-04-2002
WO 0185707 A	15-11-2001	AU 6591401 A	20-11-2001
		BR 0110703 A	28-01-2003
		CA 2407763 A1	15-11-2001

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 03/37089

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0185707 A		CN 1427830 T	02-07-2003
		WO 0185707 A1	15-11-2001
		EP 1283830 A1	19-02-2003
		JP 2003532719 T	05-11-2003
		US 2003060625 A1	27-03-2003
		US 2001051731 A1	13-12-2001
WO 0248106 A	20-06-2002	AU 3841502 A	24-06-2002
		BR 0116169 A	23-12-2003
		CA 2430579 A1	20-06-2002
		CZ 20031882 A3	17-12-2003
		WO 0248106 A2	20-06-2002
		EP 1349856 A2	08-10-2003
		NO 20032674 A	12-06-2003
		US 2002082260 A1	27-06-2002
WO 03055482 A	10-07-2003	EP 1336607 A1	20-08-2003
		WO 03047626 A1	12-06-2003
		WO 03055482 A1	10-07-2003
		US 2003138416 A1	24-07-2003
		US 2003171411 A1	11-09-2003